
**CHANGING THE MORPHOLOGICAL PARAMETERS OF THE ZEA MAYS L. PLANT,
UNDER THE ACTION DOUBLY APPLIED BY THE GRADUATION OF THE OZONE
AND ULTRAVIOLET UV-B TYPE RADIATION QUANTITY**

¹Orodan Maria, ¹Atyim Paul, ¹Glisici Mihaela,
¹Mos Liana, ¹Morgovan Claudiu, ¹Toth Csongor, ²Pallag Annamaria, ¹Osser Gyongyi

¹“Vasile Goldiș” Western University of Arad, Romania, Faculty of Medicine, Pharmacy and Dentistry,
Address: Henri Coandă St., no. 31, Arad, 310429, Romania, e-mail: gyongyiosser@ymail.com

²University of Oradea, Faculty of Medicine and Pharmacy, Department of Pharmacy

Abstract

The scientific and technical revolution has brought upon humanity long-term changes of the environment, including the deterioration of the ozone layer, with potential repercussions on the planetary life, as a consequence of the increase in the amount of UV radiation reaching the ground.

The presence of ozone in the stratosphere is essential for the existence of living creatures on Earth. The role the ozone layer plays for the biosphere lies in its ability to effectively absorb the ultraviolet (UV) radiation.

Establishing a relationship between the ozone loss in the stratosphere and the increase in UV-B radiation levels on the ground is not easy, mainly due to the difficulties in measuring the amount of ultraviolet radiation from the environment; this amount varies, depending on a number of factors, such as: greenhouse gases, aerosols, clouds, geographical positioning etc.

The ozone is responsible for the loss of a part of the harvest, which is larger than those caused by any other air pollutants. Summer is favorable to ozone production, due to the intense sunrays, high temperatures, high atmospheric pressures etc. A chronic exposure of the plants to ozone, for an extended period of time, can lead to a decrease in growth, a smaller harvest, inhibition of photosynthesis, premature aging of the leaves, a faulty distribution of the biomass, an impairment in the multiplication processes etc. The UV-B radiations have exactly the amount of energy necessary to break the chemical bond of an oxygen atom from the ozone molecule.(11)

Keywords: UV-B radiation, ozone dosage, Zea Mays, ZP471 hybrid

Submitted: 02.06.2015

Reviewed: 30.09.2015

Accepted: 22.10.2015

1.INTRODUCTION

It is therefore noted that the UV-A radiations are absorbed only in a small amount by the ozone layer, but their energy is relatively small when compared to that of the other two ranges of ultraviolet radiation. This means that the impact they have on life is significantly reduced(3-7).

The most important protection against the harmful, high-energy UV effect is ensured by the ozone layer, from the upper part of the atmosphere.(1-2)

The resulting ozone absorbs the UV-B radiations with wavelengths below 300 nm, which are very harmful to living organisms.(15)

In this way, the UV radiations help create ozone in the upper atmosphere layers.(12)

The plant growth and development is highly influenced by various environmental factors, such as: heavy metals, salinity, heat, cold, water scarcity, ozone, UV radiation etc.(5-9)

In plants, the harmful effects of these abiotic stress factors are reflected in alterations in the physiology of the plant, causing a reduction in growth and a decrease in their bioproductivity.(13)

Generally, the flavonoids from the plants' leaves have the capacity to substantially absorb UV radiations and, to a lesser extent, visible light. (14)The anthocyanins mostly absorb the visible light. The amount of flavonoids in plants varies, depending on the species and the ontogenetic development stage; quite often, the amount of flavonoids also varies between the

epidermis and mesophyll of the same leaf.(10) The flavonoid content of the leaves and their composition are frequently altered by the ambient conditions.(16-20)

MATERIAL AND METHODS


The maize plantlets subjected to the process of acclimatization originally had a height of about 9,8 to 13,7 cm and had 1-2 leaflets.

The plantlets that were prepared, which had already been subjected to the stress of irradiation with different wavelengths, according to the previous research, were placed in the ozonisation room. The soil had been watered beforehand and the plantlets periodically humidified with a pulveriser, in order to avoid their dehydration.



In the 10th day since their placement in the ozonisation room, the survival percentage of the plantlets was determined, along with their length and the number of “spots” present on their epidermis. The mean values of the indexes, marked on the 10th day since the placement of the plantlets in the room, were compared to the data registered at the start of the acclimation process; the latter was considered referential data.

RESULTS AND DISCUSSIONS

The change in length of the *Zea Mays L, Helga* leaflets, under the influence of UV-B irradiation and the gradual treatment with varying amounts of ozone is shown in the figure below.



The influence of the ozone dosage on the number of leaflets, at different values of the ozone dosage for the *Zea Mays, ZP471* hybrid



The influence of the ozone dosage on the length of the leaflets, at different values of the ozone dosage for the *Zea Mays*, ZP471 hybrid

The average number of affected leaflets had the highest readings and the irradiation with wavelengths of 280 and 310 nm, where the ozone dosage was 0,01-0,025 ppm/ ozonisation chamber volume, which resulted in a discoloration of the leaflets.

Simultaneously, the neoplantlets' leaflets, their petiole and even their stalks exhibited their epidermis sprinkled with brown-maroon specks, both on the top and bottom of the foliar limbs, slightly more intense than in the plantlets shown above (Table 75), with a yellowish – coppered color. Taking into consideration the ozone dosage adjustment, one can notice that at an insignificant dosage, the neoplantlets' leaves showed a relative increase in size.

The number of affected leaflets was relatively lower than in the case of irradiation with 287 nm wavelengths, where the ozone dosage was insignificant and medium, and the environment (0,0055 – 0,01 ppm/ ozonisation chamber volume) and caused a less intense discoloration of the intense green. The epidermis shows fewer specks of light color for the insignificant and medium ozone dosage (0,0055 – 0,01 ppm/volume), as opposed to the higher ozone dosage (0,025 ppm/volume).

The survival rate of the *Zea Mays* plantlets – the *Helga* and *ZP471* hybrids – 10 days after the commencement of the acclimatization procedure, with various ozone doses, was 100% for the three types of graduation, for the plants that were exposed to a 287 nm wavelength irradiation, compared to the surviving plantlets which were exposed to a more intense stress (310 nm).

In their natural environment, the plants are subjected to stress factors, including UV-B radiations (280 – 320 nm) which play an important role, as an increasing number of UV-B radiations reaches Earth's surface because of the stratospheric ozone layer's depletion.

The length of the leaflets exhibits changes throughout their growth, a phenomenon which is prominent at 287 nm, 10 days after the treatment. Comparing the results obtained for the *Helga* hybrid and *ZP471*, insignificant

differences were noticed. The hybrids reacted almost the same under the influence of the various ozone dosages.

REFERENCES

1. Adam, A., Bestwick, C.S., Barna, B., Mansfield, J.W., 1995, „*Enzymes regulating the accumulation of active oxygen species during the hypersensitive reaction of bean to Pseudomonas syringae pv. Phaseolica*”. *Planta* 197, pp. 240-249.
2. Aebi M.,1984. „*Catalase in vitro*”. *Methods in Enzymology* 105: pp. 121–126.
3. Aldrich, J., R., 1988. „*Chemical ecology of the Heteroptera*” *Annu. Rev. Entomol.* 33, pp. 211-238.
4. Alscher, R. G., Erturk, N., Heath, L. S., 2002, „*Role of superoxide dismutases (SODs) in controlling oxidative stress in plants*”. *J. Exp. Biol.* 2002;531(1), pp. 1331–1341
5. Ambasht, N.K., Agrawal, M., 1995, „*Physiological responses of field grown Zea mays L. plants to enhanced UV-B radiation.*” *Biotronics* 24, pp.15–23.
6. Ambasht, N.K., Agrawal, M., 1997, „*Influence of supplemental UV-B radiation on photosynthetic characteristics of rice plants*”. *Photosynthetica* 34, pp. 401–408.
7. Ghidra, V., Botu, M., Sestraş, R., I Botu, I., 2004. „*Biodiversitate și Bioconservare*”. Cluj Napoca, Editura Academic Pres.
8. Gregson, S., Clifton, S., Roberts, R.D., 1994, „*Plants as bioindicators of natural and anthropogenically derived contamination*”, *Appl. Biochemistry, Biotechnology*, 48, pp. 15-22.
9. Han, J. Lee, Y. Yeom, K. H. Kim, Y. K. Jin, H. Kim, V. N., 2004. „*The Drosha-DGCR8 complex in primary micro RNA processing*”. *Genes Dev* 18, pp. 3016–3027.
10. Haszpra, L., Barcza, Z., Davis, K. J., and Tarczay, K., 2005, „*Long term tall tower carbon dioxide flux monitoring over an area of mixed vegetation*”, *Agricultural and Forest Meteorology*, 132, pp. 58–77.

11. Haş, I., Haş V., Căbulea, I., Grecu, C., Copândeian, A., Calborean, C., Legman, V., 2004, „Ameliorarea porumbului pentru utilizări speciale”, Probl. genet. teor. aplic., XXXVI, 1-2.
12. Hausman, J.V., Kevers, C., Gaspar, T., 1994, „Involvement of putrescine in the inductive rooting phase of poplar shoots reised in vitro”, Plant Physiology, 92, pp. 201-206
13. Hausman, J.V., Kevers, C., Gaspar, T., 1995, „Auxin-poliamine interaction in the control of the rooting inductive phase of poplar shoots in vitro”, Plant Sci.110, pp. 63-71
14. Heloir, M.C., Kevers, C., Hausman, J., F., Gaspar, T., 1996, „Changes in the concentrations of auxins and polyamines during rooting of in vitro-propagated walnut shoots”. Tree Physiol 16(5): pp. 515-519.
15. Hochholdinger, F., Zimmermann R., 2008, „Conserved and diverse mechanisms in root development”. Current Opinion in Plant Biology 11, pp. 70-74.
16. Hochholdinger, F., Ling Guo, Schnable, S. 2004, „Lateral roots affect the proteome of the primary root of maize (*Zea mays L.*)”. in Plant Molecular Biology
17. Holmes, M. G., Keiller, D. R., 2002, „Effects of pubescence and waxes on the reflectance of leaves in the ultraviolet and photosynthetic wavebands: a comparison of a range of species” Plant, Cell & Environment, pp. 85-93
18. Horneck, G., 1995, „Quantification of the biological effectiveness of environmental UV radiation” Photochemistry and Photobiology B, 31, pp. 43-49
19. MONICA PAULA Raţiu, IRINA Purcărea, Florian Popa, VICTOR LORIN Purcărea, THEODOR VALENTIN Purcărea, Dumitru Lupuleasa, Daniel Boda „Escaping the economic turn down through performing employees, creative leaders and growth driver capabilities in the Romanian pharmaceutical industry”, 2011, FARMACIA; 59 vol1 119-130
20. Houghton JT, Ding Y, Griggs DJ, Noguer M, van der Linden PJ, Dai X, Maskell K, Johnson CA, 2001, „Climate change 2001. Report of the Intergovernmental Panel on Climate Change (IPCC)”. New York: Cambridge University Press.
21. Popescu, V., 1978, „Influenţa unor factori de mediu asupra creşterii şi fructificării ardeiului gras cultivat în sere ”, Teza de Doctorat IANB, Bucureşti, pp. 127-140