

EFFECT OF MICROENCAPSULATED *LACTOBACILLUS PLANTARUM* ON THE RHEOLOGICAL AND SENSORIAL PROPERTIES OF SYNBIOTIC ICE CREAM

Siti Radhiah Omar^{1*}, Siti Nazirah Omar²

¹Faculty of Science and Technology, Universiti Sains Islam Malaysia, 71800, Bandar Baru Nilai, Negeri Sembilan, Malaysia

²Faculty of Business and Management, Universiti Teknologi Mara, Malacca City Campus, 110, Off Jalan Hang Tuah, 75300, Malacca, Malaysia

*E-mail: sitiradhiah@usim.edu.my

Abstract

Synbiotic ice creams formulated with three probiotic variations (free *L.plantarum*, encapsulated *L.plantarum*, and a mixture of commercial Bifido. Sp and *L.plantarum*) was observed for their rheology and sensory properties. Microencapsulation of *L.plantarum* isolated from fermented cacao beans via freeze drying with 6% (w/v) lactose, 6% (w/v) skim milk and 3% solution of k-carrageenan coatings has shown the highest viscoelastic properties based on high fat content due to the addition of vegetable oil during the encapsulation process. Although there was no significant differences ($p < 0.05$) in the overall acceptance attributes, the degree of sweetness for encapsulated *L. plantarum* ice cream was significantly higher ($p < 0.05$) when compared to other samples. Henceforth, the encapsulation materials and freeze drying technique did improve the viscoelastic properties and overcome the sour taste of probiotics in ice cream formulation.

Keywords: Synbiotic, ice cream, encapsulation, rheology, acceptance.

Received: 30.05.2018

Reviewed: 10.09.2018

Accepted: 28.09.2018

1. INTRODUCTION

Nowadays, functional foods that could improve and confer benefit to human health are at the greatest demand (Salmeron et al., 2009; Gadhiya et al., 2015). One of the promising functional foods that is gaining its popularity and highly demanded is Probiotics. Probiotics are defined as living microorganisms which could bestow good health when consumed adequately (FAO, 2006). Referring to Tien (2009), the article (550) in the Laws of Malaysia (2006) have stated that lactic acid bacteria (LAB) is a legitimate food and the quantity must reach a minimum value of 10^6 CFU/g if the culture is to be labeled on food products. The success of probiotics in dairy products had display a positive image among consumers (Siro, et al., 2008). Probiotics are sold as supplements and has reached more than US \$28 billion worldwide in 2015, and is expected to increase up to US \$96 billion by 2020 hence, showed an optimistic future (Trend Market Research, 2018). Some of the

common probiotics are from lactic acid bacteria; *Lactobacillus sp*, and yeast such as *Saccharomyces boulardii* (Pochapin, 2000). The species of *Lactobacillus* and *Bifidobacterium* genera are commonly used as probiotics because they are known as GRAS (Generally Recognized As Safe) and is able to prove the effectiveness of the health of the human intestine (Duary et al., 2010) and also mammals (Georgieva et al., 2009). Industrially, the consumption of probiotics foods is mainly related to functional food like fermented milk, yoghurt and fermented juice (Kalita et al., 2018).

Lactic acid bacteria are gram-positive bacteria, non-spirally, non-inspired and morphologically shaped rods and cocci. It is also anaerobic facultative, negative catalase and removes lactic acid during carbohydrate fermentation. Lactic acid bacteria are more likely to grow and multiply at pH between 4.0 - 4.5 (Bamforth, 2005). Part of sepsis lactic acid bacteria can survive at a cold room temperature of 4 ° C and as high as 45 ° C. The presence of

lactic acid bacteria after 12 hours and over is as much as (10^3 CFU⁻¹ WMg), and it is capable of producing lactic acid, acetic acid and carbon dioxide (Gálvez et al., 2007). The largest group of lactic acid bacteria is made up of genus *Lactobacillus* containing more than 50 different species (De Vries et al., 2006).

The diversity of species lactic acid bacteria that dominate the fermentation of cocoa beans is limited, only species of *Lactobacillus plantarum* and *Lactobacillus fermentum* (Camu et al., 2008). According to Ardhana and Fleet (2003), lactic acid bacteria *L. cellobiosus* was dominant at 36-48 hours, reaching a maximum population growth of 10^8 - 10^9 cfu / g after 36 hours, while *L. plantarum* was dominant at the beginning of 24 hours during the process cocoa beans fermentation in Indonesia. In Malaysia, among the lactic acid bacteria found in fermented cocoa beans are *Lactobacillus collinoides*, *Lactobacillus plantarum* and some other species (Rose, 1982).

The dairy industry have found a probiotic culture as a container to be formulated in a new functional food product (Champagne et al., 2005). Yogurt and fermented milk are the main medium of probiotic culture in addition to the introduction to new products being introduced into international markets such as ice cream, butter, mayonnaise and various types of cheese. The probiotic bacteria are capsule or powdered and included in cold minced products and fermented foods (Champagne et al., 2005). As such, there are more and more probiotic products that cause consumers to be more likely to buy the product. Various strains of *L. plantarum* were marketed as probiotics (Table 2.7). According to Schillinger et al. (2005), lactic acid bacteria probiotic is often applied for fermented milk, where milk molecules can be protective against bacteria to counteract acidity of gastric juice. Shah and Ravula (2000) report that the survival of probiotic bacteria in frozen dessert desserts can be increased by encapsulation. Hence, the technique of probiotic encapsulation is able to prolong the shelf life of frozen dairy products such as ice cream.

Synbiotic ice cream is one type of functional food with the combination of both probiotics and prebiotics. Prebiotics are also known as resistant, non-digestible fiber compounds named oligosaccharides which promote the growth of probiotics either bacteria or yeast in digestive tracts. Prebiotics performed as the food supply as well as the protective carrier for the probiotics survival in human digestive system (Chen et al., 2005). Many studies have proven that consuming probiotics could confer to various health benefits. In a study by Zago et al. (2011), probiotic *L. plantarum* has shown its capability to treat many symptoms such as diarrhea and constipation. Apart from that, it could reduce the risk of getting cancer, alleviate allergic symptom, and strengthening the immune system. (Parvez et al., 2006). However, the direct addition of probiotics in food products is detrimental as the acidic environment in human gut might lead to the reduction number of their survival as they reach the intestinal tract. Moreover, their stability in the food products is extremely low and its survival in the human digestive tract is very limited. Although probiotic applications are widespread in the formulation of cheese and fermented milk, the application of this encapsulated microorganism in ice cream formulation as a medium of probiotic culture is something new (Houmayouni et al., 2008).

Owing to that reason, microencapsulation is the best protection method for bacterial cell upon cell release into the target site (Brinques, 2011). Microencapsulation technique is defined as a consolidation of food, bacterial cells, nutrients or nutraceutical substance forming small capsules which only release the contents at a particular target as a mean to protect the materials from damage and degradation due to environmental factors (Desai and Park, 2005). As contended by Nag et al. (2011), the purpose of probiotics bacteria microencapsulation process is to maintain the stability and survival during storage, to protect against environmental stress the digestive tract, and the control the release in the colon. Among the microencapsulation techniques, freeze drying is proven to be a very effective method for

encapsulation of probiotics (Semyonov et al., 2011; Tee, 2014) since the encapsulation using high temperature could kill the good bacteria (Triphati and Giri, 2014). Correspondingly, it was elucidated in some research that the survival rate for encapsulated probiotics was found significantly higher than the percentage of free cell, could offer greater protection. (Heidebach et al., 2010; Thomas et al., 2014) and pleasant taste (Anand, 2018; Tsen, 2008). As in Thomas et al. (2014), this technique could also prolong the shelf life of frozen dairy products like ice cream (Schilinger et al., 2005). Other than that, the application of microencapsulated probiotics in ice cream formulation is quite new (Houmayouni et al., 2008). Hence, the aims of this study are to focus on the assessment of sensorial, physico-chemical and as well as rheological aspect.

2. MATERIALS AND METHODS

Isolation and microencapsulation of probiotic

Lactic acid bacteria were isolated from fermenting cocoa beans together with the pulp at day 4 of the fermentation located in Raub, Pahang, Malaysia. Then, 50g of the sample was mixed with 450ml of 0.1% peptone water. In this research, pour plate technique was applied. The sample was then diluted and spread on the deMan Rogosa and Sharpe (MRS, Oxoid, UK) agar plates for further anaerobic incubation at 37°C for 24- 48 h. The colonies that growth were found to have spherical, white and creamy colour which confirm to the morphology of lactic acid bacteria. Gram staining, microscopic observation and API 50 HCL (bioMérieux, Marcy-l'Étoile, France) system were carried out to verify the finding. Then, for single colony isolation, the strain from cultured *L.plantarum* from spreading technique was taken by loop wire and was subcultured in anaerobic jar similar to previous technique mention above. Later, the cultures were stored in MRS broth composed of 20% (w/v) glycerol at -20°C for further analysis (Foong et al., 2013).

In this research, the encapsulation technique was done by referring to Otero et al. (2007). Subculturing in the MRS broth was done twice, inoculated in 2% (v/v) 800ml MRS broth and incubated at 37°C for 24-48 h. Culture was then centrifuged at (6000 x g for 10 min at 4°C) washed twice with 0.1/100 g sterile pepton water. Then, suspended bacteria was then transferred into a solution containing 6% (w/v) lactose and 6% (w/v) skimmed milk as cryoprotective agent which were believed to protect and enhance the viability of *L. plantarum* through the process of freeze drying. Twenty-five ml of the bacterial suspension was mixed with 60 ml of 3% κ-carrageenan. A mixture of 0.1% Tween 80 and 100 ml of vegetable oil with was heated at 45°C and stirred homogeneously for 3 min. Then, both mixtures were mixed and stirred slowly for 10 min. An amount of 150 ml of 0.3M KCl was poured slowly at the side of the beaker and rested for 30 min at room temperature in order to break the emulsion. Separation of oil was done by sucking and filtering microencapsulated cells. The cells were stored overnight at -20°C and freeze dried at -40°C and 0.3 mPa for 24 h using a freeze drier (Freezone Plus 6, Labconco, USA).

Preparation of synbiotic ice cream

For ice cream preparation, the materials were purchased from Giant Hypermarket, Bandar Baru Nilai, Negeri Sembilan. The synbiotic ice cream was formulated with the addition of prebiotic and probiotics following the work of Homayouni et al., (2008) with some modifications. The formula was based on the calculation of 38-39% total solids for one liter ice cream production. About 40 percent of milk and were added before elevating the temperature up to 50°C. Then, milk powder, prebiotic oligosaccharide, stabilizers, vanillin and sugar were added together into the mixture. It was homogenized at 250 bar using a homogenizer (Company Niro Soavi Italy) just before it is being pasteurized at temperature of 80°C for 20 seconds. The mixture was then cooled to 4°C and kept for 12 h to promote the ripening process of ice cream. Later, it was

poured into the ice cream maker (Taylor Company, Italy) to obtain a smooth distribution of small, disc-shaped crystals.

The next stage of synbiotic ice cream formulation was the incorporation of probiotic bacteria in which one percent (1%) of *L. plantarum* which has been encapsulated was added into a 100 ml plastic cup with out of the prepared ice cream and labeled as A. Then, one percent (1%) of free, non-encapsulated *L. plantarum* was introduced in the ice cream as B. Also, an amount of one percent (1%) free *L. plantarum* and 4% of Bifidobacteria starter culture were added together in the ice cream and labeled as C, while ice cream without any addition of microorganism was assigned to be the control sample and labeled as D. Semi-frozen mixture was cooled freeze at -18°C . During the process of freezing, frozen product temperature must be controlled (Cruz et al., 2009). Samples were stored in the freezer room for a st o and analyzed each week. Experiments were conducted in triplicate.

Rheological analysis of ice cream

Ice cream samples were fit into a rheometer (oscillatory rheometry, Physica MCR301, Anton Parr GmbH, Graz, Austria) according to the manual. Amplitude sweep test was conducted according to Waterhouse et al. (2011) to determine the linearity of the viscosity-elastic region of ice cream at a fixed frequency of 10 rad / s with strain between 0.1 to 100%. Frequency sweep test was also carried out between 0.1 and 100 rad / s at a temperature of 0 and 20°C with a gap setting, 0 mm. Data obtained in 10 points value was explained by Rheoplus V2.66 (Anton Paar GmbH, Graz, Austria).

Sensory analysis of ice cream

The organoleptic sensory for the ice cream samples were evaluated by referring to the affective test recommended by Aminah (2000). The idea st o test the level of consumer acceptance and preference towards the food samples. A total of 30 untrained consumer panels were selected randomly among the undergraduate students and graduate of Food

Biotechnology from Faculty of Science and Technology. Assessment was carried out under fluorescent light which took place in a sensory laboratory, Faculty of Science and Technology, USIM. A total of four attributes evaluated in this assessment were aroma, taste, texture and overall acceptability. The 7-point hedonic scales range from “most disliked” to “most prefer” were given to the panelists. Samples were stored and frozen at -18°C to maintain the integrity of the ice cream samples.

Statistical analysis

Statistical analysis was conducted using SAS (version 18.00) to determine the means and standard deviation for the physico-chemical and sensory data. The ANOVA Duncan’s test was conducted to determine differences between means.

3. RESULTS AND DISCUSSIONS

Biochemical API 50 CHL *L. plantarum* identification

The isolated lactic acid bacteria from *Theobroma cacao* was morphologically confirmed by gram staining, microscope and Biochemical API 50 CHL *L. plantarum* identification. It was observed that the colony showed a spherical, shiny and creamy as explained by Johansson et al. (1998). Subsequently, they were found out to be gram-positive bacteria, with rod-shape when observed under microscope which further narrow down the taxonomy of LAB. According to Hutkins (2006), Lactobacillus species has rod shaped looks which could support this research findings. Interestingly, the API 50 CHL system also shows 99.9% probability of the colonies were from *L. plantarum* species (Figure 1). The analysis of the strips after 48 hours fermentation showed some positive tests that changed the color of the bromocresole indicator purple to the yellow color. Yellow means positive while blue means negative. The black color for tubes numbered 25 is positive (Biomerieux, 2001). Color

conversion indicates that the lactobacillus is able to carry out the fermentation process and



Figure 1: Color change of fermentation reaction on API 50 CHL medium tube

use its carbohydrates and acne (heteroside, polyalcohol and uric acid) for metabolic purposes and produce acids with acidic pH of 5.2 (Barbu and Balotescu, 2006).

Table 1 shows that there are 24 types of carbohydrates that show a positive reaction of

D-Galactose, D-Glucose, D-Fructose, D-Mannose, L-Rhamnose, D-Mannitol, Methyl- α D-Mannopyranoside, N-Acetyl Glucosamine, Amygdalin, Arbutin, Eskulin citric iron, Salicin, D-Celiobiose, D-Maltose, D-Laktose, D-Mellibiose, D-Sacarose, D-Trehalose, D-Melezitose, D-Raffinose, D-Arabitol, Gentiobiose, D-Turanose and Potassium Gluconate. For the 25th tube, the color exchange is from purple to black. This is due to iron reactions with eskulin produced in the process of hydrolysis of the eskulin (Seldin and Penido, 1986).

Rheological properties of synbiotic ice cream

Ice cream stability is influenced by rheological properties as it determines the behavior of the flow and also the taste of a product. In the Amplitude Sweep Test, the viscoelasticity of each sample of ice cream has a linear response for the strain range (0.1-1%) as shown in Figure 2. In general, G' modulus of shear storage of a substance is dependent on time, which indicates the elastic properties as well as the energy saving. In this study, there is a linear state in a certain strain range because G' is independent of time, the amplitude of the swell or the swelling pressure, and also the Applied Frequency Swings.

TABLE 1: Biochemical carbohydrate metabolism profile by *L. plantarum*

Tube	Carbohydrate	Result	Tube	Carbohydrate	Result	Tube	Carbohydrate	Result
0	Control	-	19	D-Sorbitol	-	36	Amidon(Starch)	-
1	Glycerol	-	20	Metyl- α -D-Mannopiranoside	+	37	Glycogen	-
2	Erythritol	-	21	Methyl- α D-Glucopyranoside	-	38	Xylitol	-
3	D-Arabinose	-	22	N-Acetyl Glucosamine	+	39	Gentiobiose	+
4	L-Arabinose	-	23	Amygdalin	+	40	D-Turanose	+
5	D-Rybose	-	24	Arbutin	+	41	D-Lyxose	-
6	D-Xylose	-	25	Eskulin iron citrate	+	42	D-Tagatose	-
7	L-xylose	-	26	Salicin	+	43	D-Fucose	-
8	D-Adonitol	-	27	D-Celiobiose	+	44	L-Fucose	-
9	Metil- β -D xylopyranoside	-	28	D-Maltose	+	45	D-Arabitol	+
10	D-Galactose	+	29	D-Lactose	+	46	L-Arabitol	-
11	D-Glucose	+	30	D-Mellibiose	+	47	Potassium Gluconate	+
12	D-Fructose	+	31	D-Sacarose	+	48	Potassium 2-ketogluconate	-
13	D-Mannose	+	32	D-Trehalose	+	49	Potassium 5-ketogluconate	-
14	L-Sorbose	-	33	Inulin	-			
15	L-Rhamnose	+	34	D-Melezitose	+			99.9%
16	Dulcitol	-	35	D-Raffinose	+			ID
17	Innositol	-						<i>Lactobacillus plantarum</i> 1
18	D-Mannitol	+						




Figure 2: Rheological analysis of the amplitude sweep point for encapsulated *L. plantarum* ice cream, free *L. plantarum* ice cream, ice cream mix of *L. plantarum* and Bifido and also control ice cream.

The modulation of the G' modulus pattern for all ice cream samples (control, *L. plantarum* free, *L. plantarum* encapsulated and mixtures of *L. plantarum* and *Bifido*) are the same but have different amplitude when the strain or strain value is higher. G' position is consistent with the strain range (0.1-1%). The highest reading of G' value is for *L. plantarum* capsule ice cream showing a high elastic viscosity followed by control ice cream, *L. plantarum* ice cream and *Bifido* and free *L. plantarum* ice cream.

For the Frequency Sweep Test, the relationship between G' and frequency in the range 0.1 to 100 s⁻¹ were not completely linear (Figure 3). The reading of G' modulus increased with frequency in the range 0.1-100 s⁻¹. Stability of the ice cream was influenced by the rheological properties. In general, G' or the shear storage modulus of a material was time dependent, which showed the elastic and the internal, core, energy. In this study, there was a linear behavior in the range of the particular strain

because G' was not depended on time, amplitude of the oscillating strain or pressure, and also the applied frequency. The value of G' was the highest for sample A, which showed that the sample has a high viscosity elastic component. In this study, the G' modulus of all ice cream samples exhibited were depended on the frequency range tested. In addition, the high value of the elastic properties of ice cream samples also affected the slow melting rate. According to Vélez, Ruiz, and Barbosa (1997), viscosity, elastic characteristics of ice cream mix was affected by fat globule size, the distribution of air cells, the physical properties of emulsifier, protein layer, the droplets of fat, and processing procedures such as the homogenization and storage period. The instability level of both fat and protein on the surface of the micelles influenced the elastic properties of ice cream (Adapa et al., 2000). In general, encapsulated probiotics greatly enhanced the elastic and rheological properties of ice cream.




Figure 3: G' storage Modulus Analysis as a function of frequency for ice creams; (encapsulated *L. plantarum*), (free *L. plantarum*), (*L. plantarum* and Bifido) and (control)

TABLE 2: Mean scores for sensory analysis of ice cream samples

Attribute Score	4 weeks of ice cream storage at -18°C			
	Control	<i>L. plantarum</i> (free)	<i>L. plantarum</i> (encapsulated)	<i>L. plantarum</i> + <i>Bifido</i> mix
Aroma (n=30)	4.87±1.04 ^a	4.83±1.23 ^a	4.80±1.35 ^a	4.43±1.38 ^a
Texture (n=30)	4.97±1.25 ^a	4.40±1.93 ^a	5.0±1.51 ^a	4.63±1.52 ^a
Sweetness (n=30)	4.73±1.26 ^{ab}	4.43±1.38 ^a	4.77±1.57 ^a	4.4±1.35 ^{ab}
Sourness (n=30)	5.03±1.38 ^a	4.37±1.38 ^a	5.33±1.27 ^a	5.17±1.18 ^a
Overall acceptance (n=30)	5.13±0.82 ^a	4.50±1.61 ^a	5.17±1.42 ^a	4.8±1.42 ^a

^{a-b}Different alphabet at the same column showed a significant different at (p<0.05).

Table 2 shows the mean scores of ice cream perception recorded by the 30 panelists. However, in this study, the level of ice cream synbiotic acceptance was not significant at p<0.05 except for the sweetness attribute. Results found that sample A (encapsulated probiotics) showed good acceptance by sensory panels. Similar findings was reported by Nousia et al., (2011) on the acceptance of probiotics ice-cream samples incorporated with encapsulated *L. acidophilus*. Panelists thought that the sweetness is an important characteristic for an ice cream product to be acceptable. The sweet taste of ice cream mixed

with free *L. plantarum* was less favored due to higher acidity. For the encapsulated probiotics ice cream, the level of sweetness was stable and well-received by the sensory panel. According to the sensory panel, sourness was a sense of unfriendliness ice cream attribute with free probiotics. However, free *L. plantarum* ice cream was more aromatic than encapsulated *L. plantarum*. In general, the results indicate that the sample of ice cream with the assimilation of encapsulated *L. plantarum* showed higher level of optimistic acceptance among the sensory panels.

4. CONCLUSION

Encapsulated probiotics *L. plantarum* may greatly influence the rheological properties of ice cream in terms of elasticity and melting rate also could be improved by assimilating the encapsulated probiotics in the ice cream product. Ice cream prepared with freeze dried encapsulation technique may have higher viscoelasticity and slower melting rate.

Our sensory results also indicated that the ice cream made with encapsulated probiotics have higher mean scores for overall acceptability among panels which thus suitable for commercialization although it is not significant. Further shelf life study for the synbiotic ice cream and the therapeutic effects need to be conducted in the future.

ACKNOWLEDGMENTS

The authors would like to thank the authors for her technical expertise in this research project, Universiti Sains Islam Malaysia (USIM), and Universiti Teknologi Mara (UiTM), Malaysia for supporting this project. The author would like to thank to all researchers for their professional help in this research work.

5. REFERENCES

[1] Salmeron, I., Fuciños, P., Charalampopoulos, D., & Pandiella, S. S. (2009). Volatile compounds produced by the probiotic strain *Lactobacillus plantarum* NCIMB 8826 in cereal-based substrates. *Food Chemistry*, **117**(2): 265-271.

[2] Gadhiya, D., Patel, A., & Prajapati, J. B. (2015). Current Trend and Future Prospective of Functional Probiotic Milk Chocolates and Related Products--a Review. *Czech Journal of Food Science*, **33**(4).

[3] FAO (2006). Food and Agriculture Organization. 2006. Probiotics in food health and nutritional properties and guidelines for evaluation. FAO Food and Nutrition Paper. Roma: World Health Organization and Food and Agriculture Organization of the United Nations. pp. 85-112.

[4] Tien, W. H. C. (2009). Legal Status and Regulation Issues: Handbook Of Probiotics And Prebiotics, In. Lee, K. L. & Salminen, S. (Ed.), New Jersey: JohnWiley & Sons,Inc.

[5] Laws of Malaysia. (2006). Food Act 1983 and Regulation 1985. The Commissioner of Law Revision. Malaysia.

[6] Siro, I., Kapolna, E., Kapolna, B., & Lugasi, A. (2008). Functional food. Product development, marketing and consumer acceptance—A review. *Appetite*, **51**(3): 456-467.

[7] Trend Market Research. (2018). <https://www.transparencymarketresearch.com/probiotics-market.html> [Retrieved on: 18 April 2018].

[8] Pochapin, M. (2000). The effect of probiotics on *Clostridium difficile* diarrhea. *The American journal of gastroenterology*, **95**(1): S11-S13.

[9] Duary, R. K., Batish, V. K., & Grover, S. (2010). Expression of the atpD gene in probiotic *Lactobacillus plantarum* strains under in vitro acidic conditions using RT-qPCR. *Research in microbiology*, **161**(5): 399-405.

[10] Georgieva, R., Iliev, I., Haertlé, T., Chobert, J. M., Ivanova, I., & Danova, S. (2009). Technological properties of candidate probiotic *Lactobacillus plantarum* strains. *International Dairy Journal*, **19**(11): 696-702.

[11] Kalita, D., Saikia, S., Gautam, G., Mukhopadhyay, R., & Mahanta, C. L. (2018). Characteristics of synbiotic spray dried powder of litchi juice with *Lactobacillus plantarum* and different carrier materials. *LWT-Food Science and Technology*, **87**: 351-360.

[12] Bamforth, C. W. (2008). Food, fermentation and micro-organisms. John Wiley & Sons.

[13] Gálvez, A., Abriouel, H., López, R. L., & Omar, N. B. (2007). Bacteriocin-based strategies for food biopreservation. *International journal of food microbiology*, **120**(1-2): 51-70.

[14] De Vries, M. C., Vaughan, E. E., Kleerebezem, M., & de Vos, W. M. (2006). *Lactobacillus plantarum*—survival, functional and potential probiotic properties in the human intestinal tract. *International Dairy Journal*, **16**(9): 1018-1028.

[15] Camu, N., Winter, D.T., Verbrugge, K., Cleenwerck, I., Vandamme, P. & Takrama J.S. (2007). Dynamics and biodiversity of populations of lactic acid bacteria and acetic acid bacteria involved in spontaneous heap fermentation of cocoa beans in Ghana. *Applied Environment Microbiology*, **73**: 1809–1824.

[16] Ardhana, M. M., & Fleet, G. H. (2003). The microbial ecology of cocoa bean fermentations in Indonesia. *International Journal of Food Microbiology*, **86**(1): 87-99.

[17] Rose, A.H. (1982). *Economic Microbiology, Fermented Foods*, **7**. London: Academic Press.

[18] Champagne, C. P., Gardner, N. J., & Roy, D. (2005). Challenges in the addition of probiotic cultures to foods. *Critical reviews in food science and nutrition*, **45**(1): 61-84.

[19] Schillinger, U., Guigas, C., Holzapfel, W.H., (2005). In vitro adherence and other properties of lactobacilli used in probiotic yoghurt-like products. *Int. Dairy Journal*, **15**: 1289–1296.

- [20] Shah, N. P., & Ravula, R. R. (2000). Microencapsulation of probiotic bacteria and their survival in frozen fermented dairy desserts. *Australian Journal of Dairy Technology*, **55**(3): 139.
- [21] Zago, M., Fornasari, M. E., Carminati, D., Burns, P., Suárez, V., Vinderola, G., & Giraffa, G. (2011). Characterization and probiotic potential of *Lactobacillus plantarum* strains isolated from cheeses. *Food Microbiology*, **28**(5): 1033-1040.
- [22] Parvez, S., Malik, K. A., Ah Kang, S., & Kim, H. Y. (2006). Probiotics and their fermented food products are beneficial for health. *Journal of Applied Microbiology*, **100**(6): 1171-1185.
- [23] Homayouni, A., Azizi, A., Ehsani, M.R., Yarmand, M.S. & Razavi, S.H. (2008). Effect of microencapsulation and resistant starch on the probiotic survival and sensory properties of synbiotic ice cream. *Food Chemistry*, **111**: 50-55.
- [24] Brinques, G. B., & Ayub, M. A. Z. (2011). Effect of microencapsulation on survival of *Lactobacillus plantarum* in simulated gastrointestinal conditions, refrigeration, and yogurt. *Journal of Food Engineering*, **103**: 123-128.
- [25] Desai, K. G. H. & Park, H. J. (2005). Recent developments in microencapsulation of food ingredients. *Drying Technology*, **23**: 1361-1394.
- [26] Nag, A., Han, K. S. & Singh, H. (2011). Microencapsulation of probiotic bacteria using pH-induced gelation of sodium caseinate and gellan gum. *International Dairy Journal*, **21**: 247-253.
- [27] Semyonov, D., Ramon, O., & Shimoni, E. (2011). Using ultrasonic vacuum spray dryer to produce highly viable dry probiotics. *LWT-Food Science and Technology*, **44**(9): 1844-1852.
- [28] Tee, W. F., Nazaruddin, R., Tan, Y. N., & Ayob, M. K. (2014). Effects of encapsulation on the viability of potential probiotic *Lactobacillus plantarum* exposed to high acidity condition and presence of bile salts. *Food Science and Technology International*, **20**(6): 399-404.
- [29] Heidebach, T., Först, P., & Kulozik, U. (2010). Influence of casein-based microencapsulation on freeze-drying and storage of probiotic cells. *Journal of Food Engineering*, **98**(3): 309-316.
- [30] Thomas, M. B., Vaidyanathan, M., Radhakrishnan, K., & Raichur, A. M. (2014). Enhanced viability of probiotic *Saccharomyces boulardii* encapsulated by layer-by-layer approach in pH responsive chitosan-dextran sulfate polyelectrolytes. *Journal of Food Engineering*, **136**: 1-8.
- [31] Anand, S., Beniwal, A., Singh, K. S., & Aggarwal, D. (2018). Significance of probiotic encapsulation and deficiencies within.
- [32] Foong, Y. J., Lee, S. T., Ramli, N., Tan, Y. N., & Ayob, M. K. (2013). Incorporation of potential probiotic *Lactobacillus plantarum* isolated from fermented cocoa beans into dark chocolate: bacterial viability and physicochemical properties analysis. *Journal of Food Quality*, **36**(3): 164-171.
- [33] Otero, M. C., Espeche, M. C., & Nader-Macias, M. E. (2007). Optimization of the freeze-drying media and survival throughout storage of freeze-dried *Lactobacillus gasseri* and *Lactobacillus delbrueckii subsp. delbrueckii* for veterinarian probiotic applications. *Process Biochemistry*, **42**(10): 1406-141.
- [34] Cruz, A. G., Antunes, A. E. C., Sousa, A. L. O. P., Faria, J. A. F. & Saad, S. M. I. (2009). Ice cream as a probiotic food carrier. *Food Research International*, **42**: 1233-1239.
- [35] Waterhouse, D. S., Edmonds, L., Wadhwa, S. S. & Wibisono, R. (2011). Producing ice cream using a substantial amount of juice from kiwifruit with green, gold or red flesh. *Food Research International*, **5**: 30.
- [36] Aminah, A. (2000). Prinsip Penilaian Sensori. Bangi: Penerbitan Universiti Kebangsaan Malaysia.
- [37] Johansson, M. L., Nobaek, S., Berggren, A., Nyman, M., Björck, I., Ahrne, S., & Molin, G. (1998). Survival of *Lactobacillus plantarum* DSM 9843 (299v), and effect on the short-chain fatty acid content of feces after ingestion of a rose-hip drink with fermented oats. *International Journal of Food Microbiology*, **42**(1-2): 29-38.
- [38] Hutkins, R. W. (2006). Cultured dairy products. *Microbiology and Technology of Fermented Foods*, **107**.
- [39] BioMerieux. (2001). API 50 CHL medium. <http://www.biomerieux-industry.com/food/api-id32-microbial-identification>. [Retrieved on: 18 April 2018].
- [40] Barbu, V., & Balotescu, C. (2006). In vitro and in vivo studies about probiotic character of *Lactobacillus brevis* 16GAL strain. *Acta Universitatis Cibiniensis Series E: Food Technology*, **10**(1).
- [41] Seldin, L., & Penido, E. G. (1986). Identification of *Bacillus azotofixans* using API tests. *Antonie van Leeuwenhoek*, **52**(5): 403-409.
- [42] Vélez Ruiz, J. F., Barbosa Cánovas, G. V., & Peleg, M. (1997). Rheological properties of selected dairy products. *Critical Reviews in Food Science & Nutrition*, **37**(4): 311-359.
- [43] Adapa, S., Dingeldein, H., Schmidt, K. A., & Herald, T. J. (2000). Rheological properties of ice cream mixes and frozen ice creams containing fat and fat replacers. *Journal of Dairy Science*, **83**(10): 2224-2229.
- [44] Nousia, F. G., Androulakis, P. I., & Fletouris, D. J. (2011). Survival of *Lactobacillus acidophilus* LMGP.21381 in probiotic ice cream and its influence on sensory acceptability. *International Journal of Dairy Technology*, **64**(1): 130-136.