

NUTRITIONAL COMPOSITION AND BIO-EFFICACY OF ACHA (*DIGITARIA EXILIS* & *DIGITARIA IBURUA*) AND SOYBEAN (*GLYCINE MAX*) BASED COMPLEMENTARY FOODS IN RATS

Sudik, Samuel David¹, Ijarotimi, Oluwole Steve²,
Agbede, Johnson Oluwasola³ and Igbasan, Francis Adegbaye³

¹Department of Health and Animal Production, College of Agriculture, Garkawa, Plateau State, Nigeria

²Department of Food Science and Technology, Federal University of Technology, Akure, Nigeria

³Department of Animal Production and Health, Federal University of Technology, Akure, Ondo State, Nigeria

soijarotimi@gmail.com or soijarotimi@futa.edu.ng

Abstract

The nutritional potentials of acha as alternative energy source to maize in complementary foods was evaluated with a view to increase the accessibility and affordability of complementary foods by the resource poor mothers. The formulated acha-based complementary foods (*D. iburua* (DIF); *D. exilis* (DEF)), maize-based complementary food (MCF) and commercial formula (CFF) were evaluated for chemical compositions and nutritional qualities. Data were statistically analysed. Crude protein (g/100g) and energy values (kcal.) of DEF and DIF were 15.28 - 15.69 and 409.6 - 405.3 respectively, and were comparable to MCF (15.40; 407.9) and commercial formula (CFF) (15.50; 415.0). Essential amino acids of DEF and DIF were 35.64 and 38.76 g/16N respectively. The Na/K and Ca/P molar ratios of DEF and DIF ranged from 0.59 – 0.60 and 1.67 – 1.71 respectively. Phytate, oxalate and tannin concentrations were within tolerable levels. Biological values and protein efficiency ratios of experimental foods were 87.21 - 87.81% and 3.12 - 2.94 respectively, and were higher than maize-based formula (84.32%; 2.02) and commercial formula (86.53%; 2.52). Growth performance of rats fed on experimental CF were comparable to those fed on commercial formula, but lower in production costs (USD = 0.24) than CFF (USD = 0.90). The acha-based complementary foods were characterized with essential nutrients, ability to support growth and development, affordable and accessible to poor resource mothers; hence, the food samples may be suitable as complementary foods.

Keywords: Acha-based infant foods, Amino acid profiles, Nutritional qualities, Growth performance.

Received: 04.01.2019

Reviewed: 08.03.2019

Accepted: 13.03.2019

INTRODUCTION

Adequate nutrition during infancy and early childhood is essential for healthy growth and mental development of children (Kandala *et al.* 2011). Poverty and poor child feeding practices have been implicated as the major causes of protein-energy malnutrition among children in developing countries (Hanif 2011). For instance, available evidences have shown that most nursing mothers in developing countries used un-supplemented cereal porridges made from maize, sorghum and millet as complementary foods due to their inability to afford high cost fortified nutritious proprietary complementary foods (Bruyeron *et al.* 2010; Muhimbula, Issa-Zacharia & Kinabo 2011). The negative effects of this on the children however, cannot be underscored.

Recently, several efforts have been made to improve on local complementary foods through the combinations of cereals and or legumes, and other alternative locally available food resources. Such blends have been found to improve nutrient density and quality of the complementary foods with a concomitant reduction in malnutrition problems in pre-children (Beruk, Kebede & Esayas 2015; Achidi *et al.* 2016). However, information on the use of acha as alternative energy source of maize in complementary food formulation is scanty.

Acha (*Digitaria exilis* & *Digitaria iburua*), also known as fundi, hungry rice and Asian millet, is an underutilized and nutritious cereal mainly consumed as a whole grain, porridge and couscous, ground or mixed with other flours to make pastries by Africans and Asians

(Ayo & Nkama 2006; Chukwu & Abdul-kadir 2008). Acha protein is high in protein, leucine, methionine and valine than other cereals (wheat, maize, millet, sorghum, barley, oat, etc.) (Princewill-Ogbonna 2015), and these nutritional potentials appeared not to have been harnessed in the formulation of complementary foods preparations. This gap therefore forms the major focus of this present study. The study therefore aimed to formulate complementary foods from acha supplemented with soybeans flour, and also, to provide information on the chemical compositions and nutritional efficacy of the formulated products.

MATERIALS AND METHODS

Processing and Formulation of Food Samples

Acha (*Digitaria exilis* & *Digitaria iburua*), maize kernels, soybean grains, cassava starch and commercial formula were obtained from local markets in Jos, Nigeria. Each of the raw food materials was sorted, cleaned with double distilled water, oven dried at 60 °C (Plus11 Sanyo Gallenkamp PLC, UK) for 12 h, milled using Phillips laboratory blender (HR2811 model), sieved with 60 mm wire mesh (British standard) to obtain acha, maize and soybean flours respectively (Amankwah *et al.* 2009; Famakin *et al.* 2016). The experimental flour samples were blended in different proportions with reference to WHO (1994) recommended daily allowances (RDA) of energy and protein for 6 to 12 months old infant using Nutri-

Survey Linear Programming software version 2007 (Table 1).

CHEMICAL ANALYSIS

Proximate Compositions Determination

Proximate compositions (moisture content, ash, crude fiber, crude fat and crude protein content) of experimental complementary food samples were determined using the standard methods (AOAC 2012). Carbohydrate content was determined by difference as thus

$$\text{Carbohydrate (\%)} = 100 - (\% \text{Moisture} + \% \text{Fat} + \% \text{Ash} + \% \text{Crude fibre} + \% \text{Crude protein})$$

The gross energy values of the samples were determined (MJ/kg) by using Gallenkamp Adiabatic bomb calorimeter (Model CBB-330-01041; UK).

Mineral Compositions Determination

Calcium, magnesium, iron, copper and zinc were determined using Atomic Absorption Spectrophotometer (AAS Model SP9). Sodium and potassium were determined using flame emission photometer (Sherwood Flame Photometer 410, Sherwood Scientific Ltd. Cambridge, UK) with NaCl and KCl as the standards (AOAC 2012). Phosphorus was determined as phosphate by the vanadium phosphomolybdate (vanadate) colorimetric method reported by Atasié, Akinhanmi and Ojiodu (2009).

The Na/K, Ca/P, Ca/Mg, [K/Ca+Mg], Phytate: Zn, Ca: Phytate and [Ca]/[Phytate]/[Zn] molar ratios were calculated as described by Ferguson *et al.* (1988).

Table 1: Gross compositions (g) of acha-based complementary foods

Ingredients	BSD	MCF	DEF	DIF
Cassava starch	92.75	-	-	-
Maize	-	75.00	0.00	0.00
Acha Flour	-	0.00	75.00	75.00
Soybean Flour	-	17.65	17.65	17.65
Soy oil	-	4.00	4.00	4.00
Dicalcium phosphate	-	3.00	3.00	3.00
Vitamin Premix	2.45	0.25	0.25	0.25
Salt	0.3	0.10	0.10	0.10
Bine meal	4.5	-	-	-
Total	100	100.00	100.00	100.00

DIF = *D. iburua* food, DEF = *D. exilis* food, NFE= nitrogen free extract. CFF = Commercial infant foods, MCF = Maize-based complementary food, BSD= Basal diet.

Determination of Amino acid Compositions

The experimental samples were digested using 6N HCl for 24 h. Amino acids were determined using the Beckman Amino Acid Analyzer (model 6300; Beckman Coulter Inc., Fullerton, Calif., USA) employing sodium citrate buffers as step gradients with the cation exchange post-column ninhydrin derivatization method (Siswoyo *et al.* 2011). The cysteine and methionine contents were determined after performic acid oxidation and tryptophan was determined as described by Pianesso *et al.* (2015).

Determination of Anti-nutritional Factors

Phytate and tannin were determined according to the method described by Phillipson (2000, pp. 237 -48). The oxalate content of the samples was determined using the method described by AOAC (2005).

Nutritional Qualities of Experimental Food Samples in Wistar Albino Rats

Experimental design

The experimental layout was of completely randomized design (CRD). Eighty (40 males and 40 females) clinically healthy weanling Wistar albino rats (3 weeks old) were purchased from the Research rat colony, Federal University of Technology, Akure, Nigeria. The rats were divided into eight groups of 10 rats on the basis of initial weight and sex, and were individually housed in a metabolism cage (25 cm x 12 cm x 10 cm) locally designed with sufficient facilities to collect faeces and urine. The rats were fed on experimental food samples and water *ad libitum* for 28 days, weight changes and total food intake were measured. The collected faeces were oven dried at 60 °C, while urine voided was preserved in 10 mL of 10% sulfuric acid to eliminate microbial activities and prevention of nitrogen via ammonia evaporation, prior to nitrogen determination (Agbede and Aletor 2003). The feed efficiency, protein efficiency ratio, biological value and feed conversion ratio were computed (Pirman, Mari and Orenik 2007).

Biochemical Determination of Experimental Food Samples in Wistar Albino Rats

Blood and serum collection

At the end of experimental period (28 days), blood samples were collected following overnight fasting by cardiac puncture from chloroform anaesthetized rats into heparinised and non-heparinised tubes. The non-heparinised tubes were allowed to clot and were centrifuged at 3000 *xg* for 25 min to obtain the sera and stored in a deep freezer (-20 °C) prior to biochemical analysis (Agbede and Aletor 2003).

Haematological evaluation

Packed cell volume was estimated by centrifuging about 75 Fl of each blood sample in heparinized capillary tubes in a haematocrit micro-centrifuge 3000 *xg* for 5 min. Haematological indices like red blood cells, white blood cells, platelets, and haemoglobin concentration were determined (Benson and Cales 1992). The mean corpuscular haemoglobin concentration, mean corpuscular haemoglobin, and mean corpuscular volume were calculated.

Biochemical Determination

Total blood protein, creatinine and serum albumin level concentration were estimated using Randox Diagnostic Kit (Pinnell and Northam, 1978). Globulin level was determined by taking the difference between total protein and albumin (Omodanisi *et al.*, 2017). Plasma levels of alanine aminotransferase and aspartate aminotransferase activities were assayed (Varley *et al.*, 1980), while cholesterol was determined using the methods described by Wieland and Seidel (1983).

Relative organ measurements

The heart, lungs, spleen, kidneys and liver were incised, blotted free of blood, weighed and subsequently expressed in g kg⁻¹ body weight.

Ethical Approval

The study protocol was approved by the Animal Production and Health Ethical Committee of School of Agriculture and Agricultural Technology, Federal University of Technology, Akure, Nigeria with reference number: FUTA/APH-SAAT/2018/013. The

experiments on animals were conducted in accordance with the force laws and regulations as regards care and use of laboratory animals.

STATISTICAL ANALYSIS

All data were expressed as mean \pm standard error of mean (SEM) using the statistical analysis programme for social sciences (SPSS) (Edogbanya *et al.* 2016). Mean values were considered statistically significant at $P < 0.05$. Graphs were plotted using GraphPad Prism 6. Results were considered to be significant at $p \leq 0.05$.

RESULTS AND DISCUSSION

Nutrient (proximate, amino acids and minerals) Compositions of Complementary Foods

The proximate (g/100g) and energy values (kcal/100g) of acha (*D. exilis* and *D. iburua*) based complementary foods are presented in Table 2. The moisture content of the formulated experimental food samples were 4.42 and 5.45 in DEF and DIF, respectively, and these values were comparable to the value obtained for maize-based complementary food (MCF) (5.15), higher than CFF (a commercial

food) (2.00), but within the range value of FAO/WHO (1991) recommended value. The crude protein contents of DEF and DIF samples were 15.28 and 15.69, respectively, and were similar to the values obtained for maize-based complementary food (15.40), CFF (a commercial formula) (15.50) and FAO/WHO (1991) recommendation (>15). The energy values of the experimental foods were 409.6 and 405.3 in DEF and DIF respectively, and were comparable to that of MCF (407.9) and CFF (415.0), but lower than FAO/WHO (1991) recommended value (483.9). However, the energy values of these experimental complementary foods are desirable, because hundred grammes (100g) of the flour samples may provide over 80% of daily energy requirements of young children (Belscak *et al.* 2009; Lettieri-Barbato *et al.* 2012). Comparatively, the protein and energy values of acha-based complementary foods in this study were either similar or higher than what obtained by other researchers for complementary foods and cookies formulated from acha and legumes (Anuonye *et al.* 2010; Olapade and Aworh 2012; Dabels *et al.* 2016).

Table 2. Proximate compositions (g/100g) and energy values (kcal/100g) of acha (*D. exilis* and *D. iburua*) and formulated complementary foods

Parameters	WDE	WDI	MCF	DEF	DIF	CFF	RV
Moisture	12.57 $\pm 0.12^b$	13.14 $\pm 0.18^a$	5.15 $\pm 0.05^d$	4.42 $\pm 0.04^c$	5.45 $\pm 0.05^c$	2.00 $\pm 0.02^f$	<5
Total ash	5.25 $\pm 0.05^b$	5.77 $\pm 0.06^a$	2.47 $\pm 0.02^{cd}$	2.39 $\pm 0.02^d$	2.55 $\pm 0.03^c$	2.50 $\pm 0.02^{cd}$	-
Crude protein	9.72 $\pm 0.09^b$	9.93 $\pm 0.10^b$	15.4 $\pm 0.15^a$	15.28 $\pm 0.15^a$	15.69 $\pm 0.20^a$	15.50 $\pm 0.15^a$	>15
Ether extract	4.07 $\pm 0.04^d$	4.33 $\pm 0.04^d$	10.25 $\pm 0.10^b$	10.70 $\pm 0.11^a$	10.11 $\pm 0.10^b$	7.30 $\pm 0.07^c$	10-25
Crude fibre	4.92 $\pm 0.05^a$	4.06 $\pm 0.04^b$	3.20 $\pm 0.03^c$	4.17 $\pm 0.04^b$	3.31 $\pm 0.03^c$	1.70 $\pm 0.01^d$	<5
NFE	63.47 $\pm 0.63^b$	62.77 $\pm 0.60^b$	63.53 $\pm 0.81^b$	63.04 $\pm 1.20^b$	62.89 $\pm 0.71^b$	71.00 $\pm 0.71^a$	64
Energy	139.9 $\pm 1.39^b$	140.4 $\pm 1.43^b$	407.9 $\pm 4.40^a$	409.6 $\pm 4.09^a$	405.3 $\pm 4.05^a$	415.0 $\pm 4.15^a$	483.9

Means (\pm SEM) with different superscripts in the same row show significant difference at $P < 0.05$.

Key: WDI = Whole *D. iburua*, WDE = Whole *D. exilis*, MCF= maize based food, DIF = *D. iburua* food, DEF = *D. exilis* food, NFE= nitrogen free extract. CFF= Commercial infant foods. *RV (Recommended value) by FAO/WHO [1991].

The protein content of two varieties of acha kernels (9.72, 9.93 g/100g) were very low compared with the blends, this indicates that the consumption of acha alone may not produce the required level of protein in infants' complementary foods (Olapade and Aworh 2012). The improvement in nutritional qualities of food products from the blending of two or more food materials had been reported earlier by several studies (Ewuola et al., 2015; Tufa *et al.* 2016).

The amino acid compositions (g/16N) of acha (*D. exilis* and *D. iburua*) and formulated

complementary foods are presented in Table 3. The amino acid compositions of acha-based complementary foods showed that glutamic acid had the highest concentration ranging from 13.34 g/16N in DEF to 13.35 g/16N in DIF, while cysteine had the lowest value ranging from 1.21 g/16N in DIF to 1.31 g/16N in DEF. This finding agreed with the report that glutamic acid usually had the highest concentration of amino acids in plant-based foods (Anuonye *et al.* 2010, pp. 680–91).

Table 3: Amino acids profile (g/16N) of acha (*D. exilis* and *D. iburua*) and formulated complementary foods

Amino acids	WDE	WDI	MCF	DEF	DIF	*RDA
Total non-essential amino acids (TNEAA)						
Alanine	3.33±0.17 ^b	3.22±0.16 ^c	3.00±0.02 ^d	3.98±0.19 ^a	3.98±0.20 ^a	-
Aspartic acid	2.94±0.01 ^d	2.82±0.00 ^e	4.31±0.02 ^c	5.67±0.03 ^a	4.67±0.02 ^b	-
Glutamic acid	5.58±0.03 ^d	6.53±0.03 ^c	11.31±0.06 ^b	13.34±0.07 ^a	13.35±0.06 ^a	-
Serine	2.38±0.01 ^b	2.35±0.00 ^b	2.01±0.01 ^c	2.9±0.02 ^a	2.91±0.01 ^a	-
ΣNEAA	14.23±0.07 ^e	14.92±0.06 ^d	20.63±0.10 ^c	25.89±0.12 ^a	24.91±0.13 ^b	
Total conditionally essential amino acids (TCEAA)						
Arginine	3.63±0.01 ^e	4.22±0.02 ^d	5.11±0.02 ^c	6.04±0.03 ^b	6.56±0.03 ^a	2
Cystine	1.42±0.01 ^a	1.24±0.00 ^c	0.90±0.00 ^e	1.31±0.01 ^b	1.21±0.01 ^d	-
Glycine	3.31±0.02 ^d	3.49±0.02 ^b	3.42±0.02 ^c	3.84±0.01 ^a	3.84±0.02 ^a	-
Proline	2.55±0.00 ^c	2.20±0.01 ^d	2.00±0.01 ^e	3.16±0.02 ^b	3.36±0.01 ^a	-
Tyrosine	2.70±0.00 ^c	2.39±0.01 ^d	2.08±0.01 ^e	3.58±0.02 ^a	2.98±0.01 ^b	-
ΣCEAA	13.61±0.06 ^b	13.54±0.04 ^b	13.51±0.06 ^b	17.93±0.08 ^a	17.95±0.07 ^a	
Total essential amino acids (TEAA)						
Histidine	1.89±0.00 ^c	1.81±0.00 ^d	2.16±0.01 ^b	2.35±0.01 ^a	2.37±0.01 ^a	1.9
Isoleucine	3.18±0.02 ^b	2.98±0.01 ^c	2.87±0.02 ^d	2.56±0.01 ^e	3.26±0.02 ^a	2.8
Leucine	8.40±0.04 ^b	9.16±0.05 ^a	8.10±0.04 ^d	9.22±0.05 ^a	9.22±0.05 ^a	6.6
Lysine	2.80±0.01 ^d	3.14±0.01 ^c	5.20±0.02 ^b	5.72±0.02 ^a	5.72±0.01 ^a	5.8
Methionine	2.57±0.01 ^b	2.03±0.01 ^d	1.50±0.00 ^e	2.50±0.01 ^c	2.82±0.00 ^a	2.2
Phenylalanine	3.78±0.02 ^c	3.78±0.02 ^c	3.96±0.00 ^b	3.34±0.01 ^d	4.14±0.02 ^a	2.8
Threonine	3.78±0.01 ^b	4.60±0.02 ^a	2.81±0.03 ^d	3.74±0.02 ^b	3.44±0.01 ^c	3.4
Tryptophan	0.60±0.00 ^c	0.45±0.00 ^d	1.58±0.01 ^b	1.86±0.01 ^a	1.86±0.01 ^a	1.1
Valine	4.17±0.02 ^d	5.02±0.02 ^b	3.38±0.01 ^e	4.35±0.02 ^c	5.95±0.02 ^a	3.5
ΣEAA	31.17±0.16 ^d	32.97±0.21 ^c	31.56±0.15 ^d	35.64±0.18 ^b	38.76±1.01 ^a	30.1

Means (±SEM) with different superscripts in the same row show significant difference at $P < 0.05$.

Key: WDI = Whole *D. iburua*, WDE = Whole *D. exilis*, MCF = maize-based food, DIF = *D. iburua* food, DEF = *D. exilis* food, *RV (Recommended value) by FAO/WHO [1991].

The range values of total non-essential amino acids (TNEAA), conditionally essential amino acids (TCEAA) and total essential amino acids (TEAA) of acha-based complementary foods were 24.91 - 25.89, 17.93 - 17.95 and 35.64 - 38.76 g/16N, respectively. The TEAA compositions of the formulated complementary foods were higher than FAO/WHO (1991) recommended values for infants. Comparatively, the amino acid compositions of formulated foods in this study were similar to the report of other studies (Anuonye *et al.* 2010, pp. 680–91). The histidine (2.35 - 2.37 g/16N) and arginine (6.56 - 6.04 g/16N) in acha-based complementary foods were higher than FOA/WHO recommendations (1.9 and 2.0 g/day) for infants. Histidine and arginine are essential amino acids for infants, because the

guts (intestines) of infants cannot synthesized these amino acids. Besides, studies have shown that inadequate intake of arginine has been associated with many neonatal diseases, including persistent pulmonary hypertension (Vosatka *et al.*, 1994).

Minerals (mg/100g) and molar ratios of acha-based formulated complementary foods are shown in Table 4. The macro-element compositions of the acha-based complementary foods showed that magnesium had the lowest concentration, while calcium had the highest concentration with ranged values of 62.01 - 78.05 and 769.29 - 774.29, respectively. The micro-mineral elements had zinc as the lowest, while copper had the highest values with ranged values from 6.11 - 6.21 and 188.81 - 200.23, respectively.

Table 4. Minerals (mg/100g) and mineral ratios of acha (*D. exilis* and *D. iburua*) and formulated complementary foods

Parameters	WDE	WDI	MCF	DEF	DIF	*RV
Macro-elements						
Calcium (Ca)	43.30±0.22 ^c	42.36±0.21 ^c	762.00±5.38 ^b	769.29±4.61 ^{ab}	774.29±3.87 ^a	500.00
Phosphorus (P)	52.12±0.26 ^b	54.61±0.27 ^b	454.94±2.27 ^a	457.13±2.74 ^a	452.13±2.26 ^a	456.00
Potassium (K)	45.09±0.22 ^c	45.10±0.25 ^c	515.61±2.57 ^b	519.19±3.11 ^b	530.11±2.65 ^a	516.00
Sodium (Na)	37.06±0.18 ^d	37.16±0.19 ^d	302.01±1.51 ^c	335.11±2.01 ^a	319.58±1.59 ^b	296.00
Magnesium (Mg)	42.09±0.21 ^c	43.08±0.22 ^c	79.17±0.39 ^a	62.01±0.37 ^b	78.05±0.39 ^a	76.00
Micro-elements						
Iron	1.01±0.00 ^c	1.08±0.01 ^c	21.12±0.11 ^b	22.03±0.13 ^a	22.03±0.11 ^a	16
Zinc	1.56±0.01 ^c	1.73±0.01 ^b	6.18±0.03 ^a	6.11±0.04 ^a	6.21±0.03 ^a	3.2
Manganese	0.34±0.00 ^c	0.34±0.00 ^c	28.12±0.14 ^b	30.56±0.18 ^a	30.56±0.15 ^a	32
Copper	0.06±0.00 ^c	0.09±0.00 ^c	189.39±0.94 ^b	200.23±1.20 ^a	188.81±0.91 ^b	160
Mineral ratios						
[K/(Ca+Mg)]	0.53±0.00 ^c	0.54±0.00 ^c	73.85±0.36 ^a	62.68±0.37 ^b	62.68±0.31 ^b	-
Na/K	0.82±0.00 ^a	0.81±0.00 ^a	0.59±0.00 ^c	0.65±0.00 ^b	0.60±0.00 ^c	-
Ca/P	0.83±0.00 ^c	0.78±0.00 ^d	1.67±0.01 ^b	1.68±0.01 ^b	1.71±0.01 ^a	-
Ca/Mg	1.03±0.00 ^d	0.98±0.00 ^d	10.41±0.05 ^c	12.41±0.07 ^a	11.38±0.05 ^b	-

Means (±SEM) with different superscripts in the same row show significant difference at $P < 0.05$.

Key: WDI = Whole *D. iburua*, WDE = Whole *D. exilis*, MCF = maize based food, DIF = *D. iburua* food, DEF = *D. exilis* food, *RV (Recommended value) by FAO/WHO [1991].

Table 5. Antinutrient compositions (mg/100g) and phytate / (zinc & calcium) molar ratios (mol/kg) of acha (*D. exilis* and *D. iburua*) and formulated complementary foods

Parameters	WDE	WDI	MCF	DEF	DIF
Anti-nutritional factors					
Phytate	42.24±0.21 ^a	42.24±0.11 ^a	1.08±0.01 ^b	1.10±0.00 ^b	1.04±0.01 ^b
oxalate	10.10±0.05 ^a	10.20±0.05 ^a	0.16±0.00 ^b	0.17±0.00 ^b	0.18±0.01 ^b
Tannin	11.02±0.06 ^a	11.05±0.05 ^a	0.55±0.00 ^b	0.53±0.01 ^b	0.58±0.00 ^b
Molar ratios					
Ca: Phytate	0.06±0.00 ^a	0.06±0.00 ^a	0.02±0.00 ^b	0.01±0.00 ^c	0.01±0.00 ^c
Phytate: Zn	2.68±0.00 ^a	2.42±0.00 ^b	0.31±0.01 ^d	0.04±0.00 ^e	0.32±0.01 ^c
[Ca]/[Phytate]/[Zn]	2.90±0.01 ^a	2.56±0.01 ^b	0.11±0.00 ^c	0.13±0.00 ^c	0.11±0.00 ^c

Means (±SEM) with different superscripts in the same row show significant difference at $P < 0.05$.

Key: WDI = Whole *D. iburua*, WDE = Whole *D. exilis*, MCF = maize-based food, DIF = *D. iburua* food, DEF = *D. exilis* food.

Comparatively, the mineral compositions of acha-based complementary foods were higher when compared with other reports (Anigo *et al.* 2010; Ijarotimi and Oluwalana 2013). The Na/K and Ca/P molar ratios ranged from 0.59 - 0.60 and 1.67 - 1.71, respectively. It was observed in this study that all formulated complementary foods met the recommended values for Na/K (<1.0), but did not for the Ca/P (>2.0) (FAO/WHO, 1991). This observation implies that the acha-based complementary foods may not affect the regulation of heart activities of the children, however, the food samples need to be complemented with calcium for the building and maintaining of healthy bones of the children. Nutritionally, the acha-based complementary foods contain appreciable amount of essential nutrients, hence, the food samples may be suitable as a substitute for the traditional complementary food (*Ogi*, a corn gruel), which is low in vital nutrients (Anigo *et al.* 2010).

Anti-nutrient Compositions of Acha-based Complementary Foods

Antinutrient compositions (mg/100g) and phytate / (zinc & calcium) molar ratios (mol/kg) of acha (*D. exilis* and *D. iburua*) and formulated complementary foods are shown in Table 5. The range values of phytate, oxalate and tannin in acha-based complementary foods were 1.04 - 1.10, 0.17 - 0.18 and 0.53 - 0.58, respectively, and these values were lower than values obtained for complementary foods

reported by Emmanuel-Ikpeme, Ekpeyoung and Igile (2012). The range values of Ca/phytate, phytate/Zn and (Ca)(phytate) /Zn molar ratios were 0.01 - 0.02, 0.04 - 0.32 and 0.11 - 0.13, respectively, and these values were lower than the critical values reported by Oladimeji *et al.*, (2000) and Davis and Warrington (1986) for Ca/Phy (6:1) and Phy/Zn (15:1), respectively. This finding implies that the antinutrients in acha-based complementary food may not interfere with the absorption and utilisation of protein and divalent metals in the body (Greiner and Konietzny, 2006; Habtamu, 2014).

Nutritional Properties of Acha-based Complementary Foods in Wistar Rats

The growth performance of Wistar rats fed on acha-based complementary foods was comparable to those fed on commercial formula (CFF) and MCF (a maize-based complementary food) (Figure 1). This observation implies that acha-based complementary foods were of good nutritional qualities, and that the food samples may be suitable as infant foods and for the treatment of protein-energy malnutrition.

The nutritional properties of acha-based complementary foods in Wistar rats are presented in Table 6. The biological values (BV) of experimental complementary foods were 87.21 and 87.81% for DEF and DIF, respectively, and these values were significantly ($p \leq 0.05$) higher than maize-based

complementary food (MCF) (86.53%) and commercial formula (CFF) (84.32%). Protein efficiency ratios (PER) of the food samples were 3.12 and 2.94 for DEF and DIF, respectively, and these values were also significantly ($p \leq 0.05$) higher than the values obtained for MCF (2.52) and CFF (2.02). The BV and PER of these formulations met the

FAO/WHO (1991) recommended values of 70% and 2.7, respectively. In comparing with other scientific studies, it was observed that the BV of the acha-based complementary foods were similar to those reported by other researchers (Ijarotimi and Olopade 2009; Ijarotimi and Oluwalana 2013; Laminu et al. 2014).




Figure 1. Nutritional status (Weight-For-Age) of rats fed *D. iburua*-based complementary and commercial infant food (MCF = maize-based food, DIF = *D. iburua* food, DEF = *D. exilis* food, CFF (commercial formula)

Table 6. Nutritional Properties and impacts on Wister rats fed acha-based complementary foods

Parameters	MCF	DEF	DIF	CFF
Average weight gain (g/day^{-1})	4.41 \pm 0.04 ^b	4.95 \pm 0.05 ^a	5.08 \pm 0.05 ^a	3.61 \pm 0.03 ^c
Average feed intakes (g/day^{-1})	10.31 \pm 0.10 ^a	10.38 \pm 0.11 ^a	10.45 \pm 0.10 ^a	10.39 \pm 0.13 ^a
Feed conversion ratio	2.54 \pm 0.03 ^b	2.10 \pm 0.02 ^c	2.17 \pm 0.02 ^c	3.22 \pm 0.03 ^a
Biological value	86.53 \pm 0.86 ^{ab}	87.21 \pm 1.90 ^{ab}	87.81 \pm 1.87 ^a	84.32 \pm 0.84 ^b
Food Efficiency	0.39 \pm 0.00 ^c	0.48 \pm 0.01 ^a	0.46 \pm 0.01 ^b	0.31 \pm 0.00 ^d
Protein Efficiency Ratio	2.52 \pm 0.02 ^c	3.12 \pm 0.03 ^a	2.94 \pm 0.02 ^b	2.02 \pm 0.02 ^d
*Cost of feed/100g weight gain (Naira)	23.00 \pm 0.23 ^c	37.00 \pm 0.37 ^b	41.00 \pm 0.41 ^b	325.00 \pm 3.25 ^a
Relative organ weights (g/kg^{-1} body weight)				
Heart	5.11 \pm 0.05 ^b	4.95 \pm 0.04 ^b	6.48 \pm 0.06 ^a	5.04 \pm 0.05 ^b
Lungs	10.40 \pm 0.10 ^b	10.55 \pm 0.11 ^b	13.42 \pm 0.13 ^a	10.43 \pm 0.10 ^b
Spleen	4.48 \pm 0.04 ^c	4.88 \pm 0.04 ^b	6.23 \pm 0.06 ^a	4.38 \pm 0.04 ^c
Kidneys	8.25 \pm 0.08 ^b	8.51 \pm 0.09 ^b	10.42 \pm 0.10 ^a	8.46 \pm 0.08 ^b
Liver	38.10 \pm 0.38 ^b	37.56 \pm 0.37 ^b	44.54 \pm 0.44 ^a	38.74 \pm 0.38 ^b

Means (\pm SEM) with different superscripts in the same row show significant difference at $P < 0.05$.

Key: *Cost calculated on the basis of current market price. WDI = Whole *D. iburua*, WDE = Whole *D. exilis*, MCF = maize-based Food, DIF = *D. iburua* food, DEF = *D. exilis* food, CV%= coefficient of variation [%].

The relative organ weights, that is, heart, lung, spleen, kidney and liver, of rats fed on the experimental foods were comparable to those rats fed on the commercial formula.

The production costs of acha-based complementary foods (DEF and DIF) ranged from N87.21 - 87.81 (Naira) (USD = 0.24), and were higher than maize-based complementary food (N23.00) (USD = 0.06), but lower than commercial formula (CFF) (N325.00) (USD = 0.9). This observation implies that this experimental food samples was not only adequate in nutrient to support growth and development in children, but affordable by the large resource poor mothers who cannot afford

the expensive proprietary complementary foods (Bruyeron *et al.* 2010; Muhimbula *et al.*, 2011).

Biochemical Variables of Wistar Rats Fed on Acha-based Complementary Foods

The biochemical variables of rats fed on acha-based complementary foods and commercial formula are shown in Table 7. The total blood protein, serum albumen and blood cholesterol concentration of Wistar rats fed experimental foods ranged from 38.71 – 46.90 g/100g, 26.70 – 27.47 g/100g and 11.67 – 17.39 mg/100mL, respectively, and these values were comparable to those of rats fed on MCF and CFF food samples, but higher than recommended values, except for cholesterol (MVM, 2012; Wikivet, 2013).

Table 7. Biochemical properties of acha-based complementary foods in Wistar rats

Parameters	MCF	DEF	DIF	CFF	*RV
Serum metabolites and cholesterol					
Total protein (g/100g)	35.46±0.35 ^c	38.71±0.38 ^b	46.9±0.46 ^a	35.65±0.35 ^c	5.60-7.60
Albumen (g/100g)	27.81±0.27 ^a	27.47±0.25 ^{ab}	26.7±0.26 ^b	27.81±0.28 ^a	3.80-4.80
Globulin (g/100g)	7.65±0.10 ^a	7.24±0.07 ^b	2.02±0.02 ^c	7.84±0.08 ^a	-
Albumen/globulin	3.64±0.03 ^a	2.44±0.02 ^b	1.32±0.01 ^c	3.55±0.03 ^a	-
ALT (µ/DL)	9.31±0.09 ^a	9.12±0.10 ^a	8.32±0.08 ^b	9.31±0.01 ^a	17.50-30.20
AST (µ/DL)	12.4±0.11 ^b	12.17±0.13 ^b	13.7±0.10 ^a	12.3±0.12 ^b	45.70-80.80
Cholesterol (mg/100ml)	18.68±0.18 ^a	17.69±0.17 ^b	11.39±0.11 ^c	18.67±0.18 ^a	40.0-130.0
Haematological variables					
ESR (%)	0.5±0.00 ^b	0.5±0.00 ^b	0.6±0.00 ^a	0.60±0.00 ^a	-
PCV (%)	32.20±3.10 ^c	42.07±1.08 ^a	42.20±2.23 ^a	37.00±3.00 ^b	37.6-50.60
RBC (x10 ⁶ mm ³)	4.53±0.01 ^c	9.84±0.03 ^a	5.25±0.00 ^b	4.61±0.01 ^c	6.76-9.75
WBC (x10 ³ mm ³)	1.33±0.03 ^b	1.62±0.00 ^a	1.32±0.00 ^b	1.29±0.01 ^b	6.60-12.60
Hbc (g/100 mL)	11.26±0.11 ^b	14.26±0.06 ^a	10.56±0.15 ^c	10.62±0.14 ^c	11.5-16.10
MCV (µm ³)	79.18±3.03 ^a	51.58±2.71 ^b	78.86±4.10 ^a	78.6±4.03 ^a	-
MCH (pg)	16.08±2.01 ^a	15.12±1.05 ^b	16.2±0.08 ^a	15.91±0.72 ^a	-
MCHC (%)	33.3±3.00 ^a	33.32±2.03 ^a	33.03±0.98 ^a	33.3±0.34 ^a	-
Lymphocytes (%)	63.6±4.01 ^a	60.4±2.11 ^b	64.8±3.04 ^a	64.0±1.23 ^a	4.78-9.12
Neutrophils (%)	26.6±0.43 ^a	26.6±0.09 ^a	26.5±0.51 ^a	26.5±1.21 ^a	1.77-3.38
Monocytes (%)	7.18±0.00 ^a	6.6±0.04 ^b	6.6±0.02 ^b	6.6±0.00 ^b	0.01-0.04
Eosinophil (%)	2.0 ±0.00 ^a	2.0±0.00 ^a	2.0±0.00 ^a	2.0±0.00 ^a	0.03-0.08

Means (±SEM) with different superscripts in the same row show significant difference at $P < 0.05$.

MCF - maize-based food, DIF = *D. iburua* food, DIM= *D. iburua* meal, Alanine aminotransferase (ALT), Aspartate aminotransferase (AST), ESR=erythrocyte sedimentation rate, PCV= packed cell volume, RBC = red blood cells, WBC = white blood cell, Hbc= haemoglobin concentration MCV=mean cell volume, MCH=mean cell haemoglobin, MCHC=mean cell haemoglobin concentration, CFF (commercial weaning foods).

*Rat Haematological reference ranges (MVM, 2012; Wikivet, 2013).

This observation further confirmed the protein qualities of acha-based complementary foods. The range values of haematological variables, that is, pack cell volume (PCV), red blood cells (RBC), white blood cells (WBC), haemoglobin concentration (Hbc) and lymphocytes of rats fed on acha-based complementary foods were 42.07 – 42.20%, $5.25 - 9.84 \times 10^6 \text{ mm}^3$, $1.32 - 1.62 \times 10^3 \text{ mm}^3$, 10.56 – 14.26 g/100mL and 60.4 – 64.8%, respectively, and were also within the recommended values, except for WBC (MVM, 2012; Wikivet, 2013). However, slight variation was observed between the haematological indices obtained in this present study compared to the report of Aikhuomobhogbe-Arueya and Osundahunsi (2014). These variations could be attributed to the quality of diets used, concentration of proteins, breed of the rats and methods of blood analyses. The high values of PCV, RBC and Hb concentrations that were observed in acha-based complementary foods further confirmed the nutritional qualities of the formulated food products. Scientific studies have reported that Hb, PCV, MCHC and lymphocyte concentrations depend on the nutritional quality of dietary protein intake; and that diets containing poor protein may affect haemoglobin formation (Roberts et al. 2000; Oluwole et al. 2001) and immune status (Aikhuomobhogbe and Orheruata, 2006) as these indices are the major variables needed for the determination of anaemic condition of a growing child.

CONCLUSION

The present study demonstrated that the experimental complementary foods were characterized with essential nutrients, energy, ability to support growth and affordable. Hence, it could be concluded that the acha-based complementary foods in this study could be a valuable and excellent source for low-priced complementary foods, which could provide adequate nutrients for the prevention of protein-energy malnutrition in pre-school children.

Acknowledgement

The authors thank all researchers who have contributed to the success of this work, particularly in the Department of Health and Animal Production, College of Agriculture, Garkawa, Plateau State, Nigeria, and also Department of Animal production and Health, Federal University of Technology, Akure, Nigeria.

REFERENCES

- [1] Achidi, AU, Tiencheu, IB, Tenyang, N, Womeni, HM, Moyeh, MN, Ebini, LT & Tatsinkou, F 2016. Quality Evaluation of Nine Instant Weaning Foods Formulated from Cereal, Legume, Tuber, Vegetable and Crayfish, *International Journal of Food Science & Nutrition Engineering*, Vol. 6, no. 2, pp. 21-31.
- [2] Agbede, JO & Aletor, VA 2003. Comparative evaluation of weaning foods from *Glyricidia* and *Leuceana* leaf protein concentrates and some commercial brands in Nigeria, *Journal of Science of Food & Agriculture*, vol. 8, pp. 21-30.
- [3] Aikhuomobhogbe, PU & Orheruata, AM 2006. Haematological and Blood Biochemistry Indices of West Africa Dwarf Goats Vaccinated against Peste des petits Ruminats. *African Journal of Biotechnology*, vol. 5, no. 9, pp. 743 - 748.
- [4] Aikhuomobhogbe-Arueya, GL & Osundahunsi, OF 2014. Effect of extruded soy-cocoa and corn starch based complementary food on some haematological, biochemical and histopathological parameters of rats. *African Journal of Food Science*, vol. 8, no. 8, pp.419-426.
- [5] Amankwah, EA, Barimah J, Acheampong, R, Addai, LO & Nnaji, CO 2009. Effect of Fermentation and Malting on the Viscosity of Maize-Soyabean Weaning Blends. *Pakistan Journal of Nutrition*, vol. 8, no. 10, pp. 1671-1675.
- [6] Anigo, KM, Ameh, DA, Ibrahim, S & Danbauchi, SS 2010. Nutrient composition of complementary food gruels formulated from malted cereals, soybeans and groundnut for use in North-western Nigeria. *African Journal of Food Science*, vol. 4, no. 3, pp. 65-72.
- [7] Anuonye, JC, Onuh, JO, Egwim, E & Adeyemo, SO 2010. Nutrient and antinutrient composition of extruded acha/soybean blends. *Journal of Food Processing & Preservation*, vol. 34, pp. 680–691.
- [8] AOAC. 2012. Official Methods of Analysis of Association of Official Analytical Chemists, 19th edition; AOAC International: Gaithersburg, MD, USA, 2012.
- [9] AOAC. 2005. Official Methods of Analysis In: W. Horowitz, Editor, *Official methods of analysis* (18th ed.), AOAC, Gaithersburg, MD USA (2005).
- [10] Atasie, VN, Akinhanmi, TF & Ojiodu, CC 2009. Proximate analysis and physico-chemical properties

- of groundnut (*Arachis hypogea* L.). *Pakistan Journal of Nutrition*, Vol. 8, no. 2, pp. 194-197.
- [11] Ayo, JA & Nkama, I 2006. Acha (*Digitaria exilis*). *West Africa International Journal of Food and Agriculture*, Vol. 1, no. 1, pp.129-144.
- [12] Belscak, A, Komes, D, Horzic, D, Ganic, K & Damir, KD 2009. Comparative study of commercially available cocoa products in terms of their bioactive composition. *Food Research International*, vol. 42, pp. 707-716.
- [13] Benson, JP & Cales, B 1992. "Animal anatomy and physiology," in Laboratory Text Book, Wm.C. Brown Communication, Dubuque, Iowa, USA, 1992, pp. 325–341.
- [14] Beruk, BD, Kebede, A & Esayas, K 2015. Effect of Blending Ratio and Processing Technique on Physicochemical Composition, Functional Properties and Sensory Acceptability of Quality Protein Maize (QPM) Based Complementary Food. *International Journal of Food Science & Nutrition Engineering*, vol. 5, no. 3, pp. 121-129
- [15] Bruyeron, O, Denizeau, M, Berger, J & Trèche, S 2010. Marketing complementary foods and supplements in Burkina Faso, Madagascar, and Vietnam: Lessons learned from the Nutridev. Program. *Food & Nutrition Bulletin*, vol. 31, no. 2, pp. S154-S167.
- [16] Chukwu, O & Abdul-kadir, AJ 2008. Proximate Chemical Composition of Acha (*Digitaria exilis* and *Digitaria iburua*) Grains. *Journal of Food Technology*, vol. 6, no. 5, pp. 214-216.
- [17] Dabels, N, Igbabul, BD, Amove, J & Iorliam, B 2016. Nutritional Composition, Physical and Sensory Properties of Cookies from Wheat, Acha and Mung Bean Composite Flours. *International Journal of Nutrition & Food Science*, vol. 5, no. 6, pp. 401-406.
- [18] Davis, NT & Warrington, S 1986. The phytic acid, mineral, trace element, protein and moisture content of UK Asian immigrant foods. *Human Nutrition and Applied Nutrition*, vol. 40A, pp. 49–59.
- [19] Edogbanya, OP, Abolude, DS, Adelanwa, MA & Oholi, OJ 2016. The Efficacy of the Seeds of *Adansonia digitata* L. as a Biocoagulant and Disinfectant in Water Purification. *Journal of Earth Environment and Health Science*, vol. 2, pp.122-128.
- [20] Emmanuel- Ikpeme, CA, Ekpeyoung, IO & Igile, GO 2012. Nutritional and Sensory Characteristics of an Infant Food Based on Soybean Seeds (*Glycine max*) and Tigernut Tubers (*Cyperus esculenta*). *British Journal of Applied Science & Technology*, vol. 2, no. 4, pp. 356-366.
- [21] Ewuola, GO, Ibironke, SI & Fashakin, JB 2015. Formulation and Nutritional Evaluation of Maize, Bambara Groundnut and Cowpea Seeds Blends Complementary Food. [American Journal of Food & Nutrition](#), vol. 3, no. 4, pp.101-105.
- [22] Famakin, O, Fatoyinbo, A, Ijarotimi, OS, Badejo, AA & Fagbemi, TN 2016. Assessment of nutritional quality, glycaemic index, antidiabetic and sensory properties of plantain (*Musa paradisiaca*)-based functional dough meals. *Journal of Food Science & Technology*, vol. 29, pp.1-11.
- [23] FAO/WHO. 1991. CODEX CAC/GL. Codex Alimentarius: Guidelines on Formulated Supplementary Foods for Older Infants and Young Children. Vol. 4, FAO/WHO Joint Publications: p. 144, 1991.
- [24] Ferguson, EL, Gibson, RS, Thompson, LU & Berry, M 1988. Phytate, zinc, and calcium contents of 30 East African foods and their calculated phytate: Zn, Ca: phytate, and [Ca][phytate]/[Zn] molar ratios. *Journal of Food Composition Analysis*, vol. 1, no. 4, pp. 316-325
- [25] Greiner, R. and Konietzny U. 2006. Phytase for Food Application. *Food Technology & Biotechnology*, vol. 44, pp.125-140.
- [26] Grubben, G.J.H and Denton, O.A. (2016). Vegetables. *Plant Resources of Tropical Africa*. p. 282. ISBN 978-90-5782-147-9.
- [27] Habtamu, FG 2014. Potential Health Benefits and Adverse Effects Associated with Phytate in Foods, *Food Science and Quality Management*, vol. 27, pp. 45-54.
- [28] Hailu, S, Seyoum, T and Dechassa, N (2008). Effect of combined application of organic- P and inorganic-N fertilizers on yield of carrot. *African Journal of Biotechnology*, 7 (1) 27-34
- [29] Hanif, HM 2011. Trends in breastfeeding and complementary feeding practices in Pakistan, 1990-2007. *International Breastfeeding Journal*, vol. 6, no.15, pp. 1-7.
- [30] <https://books.google.com> (Accessed June 22nd, 2018)
- [31] <https://en.climate-data.org> (2018). Climograph and Temperature graph of Ilorin
- [32] <https://en.m.wikipedia.org/wiki/Carrot>. Org /wiki /carrot. (Accessed June 26th, 2018)
- [33] <https://link.springer.com> (2017)
- [34] Ijarotimi, OS & Olopade, AJ 2009. Determination of Amino Acid Content and Protein Quality of Complementary Food Produced from Locally Available Food Materials in Ondo State, Nigeria. *Malaysian Journal of Nutrition*, vol. 15, no. 1, pp. 87-95.
- [35] Ijarotimi, OS & Oluwalana, IB 2013. Chemical Compositions and Nutritional Properties of Popcorn-Based Complementary Foods Supplemented with *Moringa oleifera* Leaves Flour, *Journal of Food Research*, vol. 2, no. 6, pp.117-132.
- [36] Kandala, N, Madungu, TP, Emina, JBO, Nzita, KPD & Cappuccio, FP 2011. "Malnutrition among Children Under the Age of Five in the Democratic Republic of Congo (DRC): Does Geographic Location Matter?," *BMC Public Health*, vol. 11, p. 261.

- [37]Kiran.M, Jilani.M. S, Waseem. K and Marwai,S.K.(2016). Response of Carrot (*Daucus carota*) Growth and Yields to Organic and Inorganic manure. *American- Eurasian Journal. Agricultural and Environmental Sciences*. 16(6): 1211- 1218.
- [38]Kochhar, S. L (2011). Economic Botany in the Tropics. Fourth Edition, pp245-246. Macmillan
- [39]Laminu, HH, Sheriff, M, Bintu, BP & Muhammad, AA 2014. Evaluation of the Protein Quality of Composite Meals produced from Selected Cereals and Legumes for infants. *Scholarly Journal of Agricultural Science*, vol. 4, no. 11, pp. 536-542.
- [40]Lettieri-Barbato, D, Villano, D, Beheydt, B, Guadagni, F, Trogh, I & Serafini, M 2012. Effect of ingestion of dark chocolates with similar lipid composition and different cocoa content on antioxidant and lipid status in healthy humans. *Food Chemistry*, vol. 132, pp. 305-1310.
- [41]Mehedi. T. A., Siddique. M. A and Sonia, B. S. (2012). Effects of Urea and Cowdung on growth and yield of Carrot. *Bangladesh Agricultural Research Institute, Gazipur. p.94. 10 (1): 9- 13*.
- [42]Muhimbula, HS, Issa-Zacharia, A & Kinabo, J 2011. Formulation and sensory evaluation of complementary foods from local, cheap and readily available cereals and legumes in Iringa, Tanzania. *African Journal of Food Science*, vol. 5, no. 1, pp. 26 - 31.
- [43]MVM 2012. Haematologic reference ranges. Merck Veterinary Manuals.com; 2012.
- [44]Oladimeji, MO, Akindahunsi, AA & Okafor, AF 2000. Investigation of the bioavailability of zinc and calcium from some tropical tubers. *Nahrung*, vol. 44, pp. 136–137.
- [45]Olapade, AA & Aworh, OC 2012. Evaluation of extruded snacks from blends of acha (*Digitaria exilis*) and cowpea (*Vigna unguiculata*) flours. *Agricultural Engineering International: CIGR Journal*, vol. 14, no. 3, pp. 210–217.
- [46]Oluwole, AK, Olorunfunmilayo, OA, Odubiyi, AO & Babatunde, GM 2001. Comparative effects of feeding dried poultry versus dried swine wastes on the haematological and serum biochemical parameters of weaner pigs in the tropics. *Tropical Animal Production Investment*, vol. 4, pp. 91–98.
- [47]Omodanisi, EI, Aboua, YG & Oguntibeju, OO 2017. Assessment of the Anti-Hyperglycaemic, Anti-Inflammatory and Antioxidant Activities of the Methanol Extract of *Moringa Oleifera* in Diabetes-Induced Nephrotoxic Male Wistar Rats. *Molecules*. Vol. 22, no. 439, pp.1-7.
- [48]Phillipson, JD 2000. Phytochemistry of medicinal plants. *Phytochemistry*, vol. 56, no. 3, pp. 237 -248.
- [49]Pianesso, D, Radünz Neto, J, da Silva LP Goulart, FR., Adorian, TJ, Mombach, PI, Loureiro, BB, Dalcin, MO, Rotili, DA & Lazzari, R 2015. Determination of tryptophan requirements for juvenile silver catfish (*Rhamdia quelen*) and its effects on growth performance, plasma and hepatic metabolites and digestive enzymes activity. *Animal Feed Science & Technology*, vol. 210, pp.172–183.
- [50]Pinnell, AE & Northam BE 1978. New automated dye-binding method for serum albumin determination with bromocresol purple. *Clinical Chemistry*. Vol. 24, pp. 80-86.
- [51]Pirman, T, Mari, M & Orenik, A 2007. “Changes in digestibility and biological value of pumpkin seed Cake protein after limiting amino acids supplementation,” *Izvorni znanstveni članak*. Vol. 49, no. 2, pp. 95–102.
- [52]Princewill-Ogbonna IL (2015). Proximate and Anti-Nutritional Studies of Two Varieties of Germinated Acha Seed (*Digitaria Exilis*). *The International Journal of Science & Technology*. Vol.3, no. 4, pp. 213-218.
- [53]Roberts, KM, Daryl, KG, Peter, AM & Victor, WR 2000. Mayer’s Biochemistry, 25th edition, McGraw Hill, New York, vol. 25, pp. 763-765.
- [54]Siswoyo, TA, Mardiana, E, Lee, KO & Hoshokawa, K 2011. Isolation and characterization of antioxidant protein fractions from melinjo (*Gnetum gnetum*) seeds. *Journal of Agriculture & Food Chemistry*, Vol. 59, pp. 5648–5656.
- [55]Tiwari, B. K., Brunton, Nigel P., Brennan, Charles (2012). *Handbook of Plant Food Phytochemicals: Sources, Stability and Extraction* . John Wiley & Sons. p. 405. ISBN 978-1-118-46467-0
- [56]Tufa, MA, Urga, K, Weledesemayat, GT & Mitiku, BG 2016. Development and Nutritional Assessment of Complementary Foods from Fermented Cereals and Soybean. *Journal of Food Science & Nutrition*, vol. 2, no. 014, pp. 1-8.
- [57]Uddain, J., Chowdhury, S and Rahman, M, J (2010). Efficacy of different organic manures on growth and productivity of Radish (*Raphanus sativus L.*) and Carrot (*Daucus carota L.*). *IAEB 3(2): 189-193*.
- [58]Varley, H, Gewenlock, A & Bell, M 1980. Practical Clinical Biochemistry, 1. 5th ed., 741, 897. London: William Heinemen Medical. Books, Ltd.
- [59]Vosatka, RJ, Kashyap, S & Trifiletti, RR 1994. Arginine deficiency accompanies persistent pulmonary hypertension of the newborn. *Biology Neonate*, vol. 66, pp. 65–70.
- [60]WHO 1994. Codex Alimentarius Standards for Foods for Special Dietary Uses (including foods for infants and children, Vol. 4. Joint FAO/WHO Food Standards Program Rome: WHO, Codex Alimentarius Commission.
- [61]Wieland, H & Seidel, D 1983. A fully enzymatic colorimetric determination of HDL-cholesterol in the serum. *Journal of Nutrition*, vol. 109, pp. 760-766.
- [62]Wikivet 2013. Haematology. Available at: en.wikipedia/wiki/haematology (accessed May, 2017).
- [63]World Sites Atlas (2018). Latitude and Longitude of Ilorin
- [64]www.fspublishers.org. (Accessed July 19th, 2018)